

CUVRIOR (trientine tetrahydrochloride) is a prescription medicine approved for adults with stable Wilson Disease (WD) who are de-coppered and tolerant to penicillamine.

What to consider before talking to your doctor

- **What treatment** are you currently taking?
- **What challenges do you have** with your medication for WD (check all that apply)
 - Frequency of dosing
 - Number of pills per dose
 - Taking your medication as prescribed
 - Keeping medication in the refrigerator
 - Food restrictions around medication
 - Difficulty swallowing
 - Tolerability of medication
 - Cost of medication
- **Have you been experiencing** any additional symptoms while on your current medication?

What to ask your doctor

- **Should I be monitoring my WD more frequently** than I currently am?
- **Are there any additional tests**, beyond 24-hour urinary copper test, to ensure my copper levels are controlled properly?
- Based on the challenges with my current therapy, **are there other medications that would make it easier to take my medication as prescribed?**
- **What would the transition plan look like** if I decided to switch to a new treatment?
- **Will my insurance allow me to switch?**
 - If a Prior Authorization is required, how would we navigate that together?



Tell your doctor if you are interested in CUVRIOR as your maintenance treatment for WD

- No refrigeration required
- Twice-daily dosing
- Scored tablets

IMPORTANT SAFETY INFORMATION

- **Do not use** CUVRIOR if you are allergic to trientine or to any of the ingredients in CUVRIOR.

Please see additional Important Safety Information on back.

Ask your doctor if CUVRIOR is right for you

For more information about CUVRIOR or patient resources that may be available to you, please visit www.CUVRIOR.com or scan the QR code.



Notes

INDICATION

CUVRIOR is a prescription medicine used to treat adults with stable Wilson Disease who are de-coppered and able to take penicillamine. Wilson Disease is a condition where the body stores too much copper. CUVRIOR is a chelating medicine which binds copper in the blood stream and eliminates it in the urine to help reduce excess copper in the organs.

IMPORTANT SAFETY INFORMATION

- **Do not use** CUVRIOR if you are allergic to trientine or to any of the ingredients in CUVRIOR.
- **Your Wilson Disease symptoms could get worse when you start treatment.** This could happen because too much copper is removed from the body in a short period of time. Your doctor may need to reduce your dose or stop CUVRIOR treatment.
- **Copper deficiency** may develop following treatment with CUVRIOR. Your doctor will do tests to monitor your urine and blood for copper.
- **Iron deficiency** may develop while taking CUVRIOR. If this happens, your doctor may tell you to take iron supplements for a limited time.
- **Allergic reactions**, such as a rash can occur. If a rash or other allergic reaction occurs, contact your doctor immediately or get emergency help.
- The most common side effects are pain in the abdomen, change in bowel habits, rash, hair loss (alopecia), and mood swings.

Take CUVRIOR at least 1 hour apart from any other oral medicine. Tell your doctor about all the medicines you're taking, including over-the-counter medicines, vitamins, and herbal supplements. Avoid taking supplements when taking CUVRIOR. Taking CUVRIOR with mineral supplements (e.g., iron, zinc, calcium, magnesium) can reduce the effectiveness of CUVRIOR. If iron supplementation is necessary, take CUVRIOR at least 2 hours before or 2 hours after taking iron; for other mineral supplements (e.g., zinc, calcium, magnesium), take CUVRIOR at least 1 hour before or 2 hours after. Do not start or stop any medicines while taking CUVRIOR without talking to your doctor.

These are not all the possible side effects of CUVRIOR. You are encouraged to report negative side effects of prescription drugs. Contact Orphalan at 1-800-961-8320 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see full **[Prescribing Information](#)**.

